



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A systematic review of behaviour change techniques within interventions to prevent return to smoking postpartum

Citation for published version:

Brown Tracey, J, Wendy, H, Bauld, L, Richard, H, Vivienne, M, Felix, N, Sophie, O, Michael, U & Caitlin, N 2018, 'A systematic review of behaviour change techniques within interventions to prevent return to smoking postpartum', *Addictive Behaviors*. <https://doi.org/10.1016/j.addbeh.2018.12.031>

Digital Object Identifier (DOI):

[10.1016/j.addbeh.2018.12.031](https://doi.org/10.1016/j.addbeh.2018.12.031)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Addictive Behaviors

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Accepted Manuscript

A systematic review of behaviour change techniques within interventions to prevent return to smoking postpartum

J. Brown Tracey, Hardeman Wendy, Bauld Linda, Holland Richard, Maskrey Vivienne, Naughton Felix, Orton Sophie, Ussher Michael, Notley Caitlin



PII: S0306-4603(18)31324-8
DOI: <https://doi.org/10.1016/j.addbeh.2018.12.031>
Reference: AB 5851
To appear in: *Addictive Behaviors*
Received date: 14 November 2018
Revised date: 13 December 2018
Accepted date: 22 December 2018

Please cite this article as: J. Brown Tracey, Hardeman Wendy, Bauld Linda, Holland Richard, Maskrey Vivienne, Naughton Felix, Orton Sophie, Ussher Michael, Notley Caitlin , A systematic review of behaviour change techniques within interventions to prevent return to smoking postpartum. *Ab* (2018), <https://doi.org/10.1016/j.addbeh.2018.12.031>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A systematic review of behaviour change techniques within interventions to prevent return to smoking postpartum

Brown Tracey J MMedSci*¹, Hardeman Wendy PhD², Bauld Linda PhD³, Holland Richard PhD⁴, Maskrey Vivienne MSc¹, Naughton Felix PhD², Orton Sophie PhD⁵, Ussher Michael PhD⁶, Notley Caitlin PhD¹

1 Norwich Medical School, University of East Anglia, Norwich, UK

2 School of Health Sciences, University of East Anglia, Norwich, UK

3 Usher Institute, College of Medicine and Veterinary Medicine, University of Edinburgh, Edinburgh UK

4 Leicester Medical School, University of Leicester, Leicester, UK

5 Division of Primary Care, University of Nottingham, Nottingham, UK

6 Population Health Research Institute, St George's, University of London, London, UK & Institute for Social Marketing, University of Stirling, Stirling, UK

* Corresponding author:

Tracey Brown: tracey.j.brown@uea.ac.uk

Abstract

Introduction

There is no routine support to prevent postpartum smoking relapse, due to lack of effective interventions. Previous reviews have identified behaviour change techniques (BCTs) within pregnancy cessation trials to specify which components might be incorporated into more effective interventions, but no reviews have identified BCTs for prevention of smoking relapse postpartum. We reviewed BCTs and potential delivery modes, to inform future interventions.

Methods

We searched Medline and EMBASE from January 2015- May 2017; and identified trials published before 2015 by handsearching systematic reviews. We included RCTs where: i) ≥ 1 intervention component aimed to maintain smoking abstinence versus a less intensive intervention; ii) participants included pregnant or postpartum smoking quitters; iii) smoking status was reported in the postpartum period. We extracted trial characteristics and used the Behaviour Change Technique Taxonomy v1 to extract BCTs. We aimed to identify 'promising' BCTs i.e. those frequently occurring and present in ≥ 2 trials that demonstrated long-term effectiveness (≥ 6 months postpartum). Data synthesis was narrative.

Results

We included 32 trials, six of which demonstrated long-term effectiveness. These six trials used self-help, mainly in conjunction with counselling, and were largely delivered remotely. We identified six BCTs as promising: 'problem solving', 'information about health

consequences', 'information about social and environmental consequences', 'social support', 'reduce negative emotions' and 'instruction on how to perform a behaviour'.

Conclusions

Future interventions to prevent postpartum smoking relapse might include these six BCTs to maximise effectiveness. Tailored self-help approaches, with/without counselling, may be favourable modes of delivery of BCTs.

Registration: PROSPERO CRD42018075677

Keywords

smoking relapse; intervention; postpartum; pregnancy; review; behaviour change techniques

Introduction

Smoking in pregnancy or postpartum remains a major preventable cause of maternal and infant mortality and morbidity.¹ Pregnancy acts as a strong motivator for smoking cessation, with more women quitting during pregnancy than at any other time in the life course. Indeed, it is estimated that up to 49% of women are able to 'spontaneously quit' before their first antenatal appointment² due to factors such as maternal concerns for the health of the unborn baby.^{3,4} Pregnant women can be considered a particularly vulnerable population, since not only are they themselves at risk of harm from tobacco smoke, but there are also increased health risks to the developing fetus and to infants born to smoking mothers. Cessation support, including the UK NHS Smokefree services and provision of carbon monoxide

monitoring, are effective and cost effective ways to support pregnant smokers to quit.⁵ Most women who quit smoking during pregnancy wish to remain abstinent, but a high proportion return to smoking postpartum.^{2,6,7} Postpartum women are particularly vulnerable to smoking relapse due to factors such as perception of no longer needing to protect the baby, stress of dealing with a new infant, lack of confidence in remaining quit, nicotine dependence, living with a smoking partner, and a desire to return to their pre-pregnancy identity; act as contributors to relapse post-birth.^{3,4,8,9} Rates of return to smoking in the literature vary due to specific population characteristics and variation in success of any provided intervention.^{6,10} Recent data estimates that up to 76% of spontaneous quitters restart smoking postpartum.⁶ For women receiving cessation support; who may be more addicted or have weaker beliefs about the dangers of smoking; rates may be higher.^{2,6,7} In randomised clinical trials of within-pregnancy smoking cessation interventions, secondary analysis using point-prevalence data estimated that across trials, the mean proportion of women smoking at the end of pregnancy was 87%, rising to 94% six months later, suggesting the majority of smoking cessation trial participants continue to smoke both throughout pregnancy and after childbirth.⁶

Return to smoking postpartum has substantial health and cost implications, thus reducing exposure to smoke in pregnancy is an important goal worldwide.¹¹ In the UK alone, the annual cost to the NHS of continuing to smoke in pregnancy is estimated to be as high as £64 million for treating health related problems for mothers; and costs for treating mortality and morbidity of infants (aged 0-12 months) due to maternal smoking an estimated further £23.5 million.¹² Sustained smoking abstinence postpartum has significant health benefits for the mother and wider family due to reduced exposure to second hand smoke. Quitting smoking before the age of 30 avoids more than 97% of the excess mortality caused by smoking, due to lower risks of cancers, coronary heart disease and cerebrovascular disease.¹³ Infants exposed

to second hand smoke have a higher incidence of sudden infant death, respiratory conditions including asthma, bronchitis and pneumonia, and other infections such as middle ear disease and meningitis.¹⁴ Furthermore, children of smoking mothers are twice as likely to become smokers themselves, perpetuating the cycle.¹⁵ The prevalence of smoking during pregnancy is higher in more disadvantaged groups^{2,16} and being less well educated has been identified as an important predictor of relapse postpartum.⁸ Despite these inequalities, health and cost implications; there is no routine provision of support to prevent relapse due to a lack of evidence-based interventions.¹⁷ There is thus a clear need to develop an effective intervention to prevent return to smoking in postpartum women.

A number of interventions for relapse prevention in pregnancy and maintenance into the postpartum period have been developed; with modest success for those incorporating counselling, health education and/or financial incentives.² Behavioural change interventions are complex in nature, consisting of multiple components designed to affect change.¹⁸ In order to describe active ingredients (components) of interventions, Michie and colleagues developed the Behaviour Change Technique Taxonomy v1 (BCTTv1).^{19,20} They propose that by identifying individual behaviour change techniques (BCTs) within interventions, it is possible to specify how interventions operate and therefore what components might be incorporated into new and more effective interventions.²⁰ Other reviews have identified BCTs for a range of behaviour change interventions including smoking cessation support for general smokers^{21,22} and for smoking cessation in pregnancy^{23,24}; but no reviews have identified BCTs for smoking relapse prevention in the postpartum period. We used the BCTTv1 taxonomy¹⁹ to identify BCT components which could be considered 'promising, or likely to be effective in future interventions to maintain smoking abstinence postpartum. We also reviewed the characteristics (mode of delivery and fidelity) of smoking relapse

prevention interventions in order to understand how these ‘promising’ BCTs might be best delivered.

Methods

Types of trials

We included randomised controlled trials of interventions where at least one component was designed to maintain smoking abstinence, versus a less intensive intervention or usual care. Participants were generally healthy pregnant or postpartum women who were recent smoking quitters (or a mixed population of recent quitters and smokers at baseline). Trials had to report smoking status (self-report or biochemically validated) in the postpartum period (beyond delivery). Trials not published in English were excluded due to the detailed intervention information required. Our review was registered on the PROSPERO international prospective register of systematic reviews (CRD42018075677).

Searches

We searched Medline and EMBASE from 1st January 2015 to 9th May 2017, to replicate and update the search strategy used by Jones and colleagues⁶ in a published systematic review of restarting smoking in the postpartum period. Keywords included: pregnancy, antenatal, prenatal, childbirth, postnatal, postpartum, breastfeeding, fetus, newborn, infant, tobacco, smoking, smoking cessation, and relapse. The full search strategy is published in Jones.⁶

Randomised controlled trials published prior to 2015 were identified by handsearching Jones⁶ and other recent pertinent systematic reviews of smoking cessation in pregnancy^{2,16} and postpartum relapse prevention interventions.^{4,8}

Selection of trials

One author (TJB) screened citations on the basis of title and abstract, and also using tables of study characteristics when handsearching. A second reviewer (IG) independently screened 10% of titles and abstracts for studies that appeared to be relevant to the review.

Disagreement rate was not formally measured, but any disagreements were resolved by consensus. Where it was unclear if a study met our inclusion criteria, the full-text was collected and assessed. Each full-text article was assessed for inclusion using an inclusion/exclusion log. For excluded studies, the reason for exclusion was recorded at full-text assessment.

Data extraction

We identified BCTs targeting smoking behaviour, i.e. BCTs included to maintain smoking abstinence, or to promote smoking cessation together with maintaining smoking abstinence. The BCTTv1 was used to extract BCTs by assigning BCT codes to appropriate sections of trial articles.¹⁹ Two researchers (TJB and CN) completed online training in BCTTv1 use (www.bct-taxonomy.com). To ensure consistency in data recording, data extraction of BCTs was duplicated independently at a level of 10%, with any disagreements resolved by discussion or the involvement of an expert third reviewer (WH) who co-authored the BCTTv1.^{19,20} To ensure any relevant BCTs were captured, we extracted BCTs whether they were definitely (coded ++) or probably (coded +) present, following BCTTv1 coding principles (www.bct-taxonomy.com). We calculated frequency of occurrence of BCTs across trials to define a list of 'promising' BCTs, most likely to be effective in relapse prevention interventions. We defined a BCT as 'promising' if it was present in at least two long-term effective interventions, as in Lorencatto²⁴, and where frequency in all included trials, regardless of trial effectiveness, was highest (present in $\geq 25\%$ interventions). We defined

long-term effective interventions as those finding statistically significant ($p < 0.05$ where reported) differences in smoking abstinence between the intervention and control groups at six months postpartum or later (biochemically validated or self-report). The rationale for this two-pronged approach, was that by considering BCTs in light of their overall frequency, we might identify techniques most likely to be feasible, acceptable and fit for purpose.

We also extracted data concerning characteristics of trial participants, details of the intervention and control, smoking status outcomes and data on intervention fidelity (adherence to the protocol, acceptability, compliance) in order to consider BCTs which might be most effective and suitable for postpartum relapse interventions and how these interventions might be best delivered. Data synthesis was narrative.

Results

Results of the search

The inclusion of RCTs is summarised in Figure 1. Electronic searching identified 636 records, 88 duplicate references were removed, resulting in 548 references for title and abstract screening. From these records, 72 relevant full text articles were identified (48 unique trials). Handsearching identified a further 69 relevant trials (after removal of duplicates). In total, articles from 117 studies were retrieved for full text assessment with the final inclusion of 32 primary RCTs (106 articles) which satisfied all inclusion criteria.²⁵⁻¹³⁰

Characteristics of all included trials

The details of participants, intervention, comparison group, outcome measures, main findings and BCTs identified are described in Supplementary Table 1. The majority of studies (25 trials) were conducted in the USA,^{25,26,34,36,57,64,66,75,83-86,88,89,94,98,99,101,106,111,113,116,118,124,128} two in Canada,^{55,80} two in the UK,^{67,129} and three elsewhere in Europe (Germany, Spain and Poland)^{68,79,95}; all notably high income countries. Less than a third of interventions targeted lower socioeconomic groups or ethnic minorities.^{36,83,84,86,106,111,118,124,128,129}

Interventions were heterogeneous, differing in intervention content, mode of delivery, and population. Thirteen trials included mixed populations of pregnant smokers and quitters^{34,36,55,67,75,84,86,94,95,98,101,111,113}; eleven included pregnant quitters^{26,57,64,66,83,99,106,116,118,124,128}; two included mixed populations of postpartum smokers and quitters^{68,79}; two included postpartum quitters^{25,80}; three included pregnant women and partners^{85,89,129}; and one included postpartum women and partners.⁸⁸ Interventions in most trials (n=21) consisted of face-to-face or telephone advice (counselling, cognitive behavioural therapy, motivational interviewing) plus additional support (mainly written self-help materials), with the remainder using: counselling alone^{36,116,118}; self-help materials alone^{26,89,94,98}; oral progesterone^{25,66}; mobile phone alerts⁶⁴; incentives plus counselling and supporting materials.⁸⁴ Interventions were delivered by a wide range of clinical staff, pregnancy specialists, counsellors or researchers in either home or clinic settings. The intensity, duration, and time between intervention sessions varied widely from the provision of a single session to a maximum of 14 sessions provided up to 9 months postpartum. Only two trials followed up participants beyond 12 months postpartum.^{68,116} We found no clear pattern as to which modes of delivery and timings of intervention contact would be best used to deliver intervention components most effectively.

Smoking status was validated in at least a sample of the population at a particular time point in 26 trials, but six trials only used self-report data.^{55,64,68,83,95,129} Many trials reported some fidelity and/or acceptability issues, and we considered 10 trials as having reported major intervention fidelity and/or acceptability concerns,^{55,67,68,75,80,83,86,101,113,129} which may have influenced intervention effectiveness. These trials highlighted issues of low recruitment, low level of intervention implementation, low level of uptake of the intervention, or high dropout rates. There was no distinct pattern as to which specific components of interventions had been most affected by fidelity and/or acceptability, since this level of detail was often not reported. Some trials made no reference, or minor reference, to fidelity related issues suggesting there may also have been some underreporting of concerns relating to protocol adherence, acceptability, or compliance.

Characteristics of long-term effective trials

We found six long-term effective trials demonstrating a statistically significant difference in smoking abstinence between the intervention and control groups at six months postpartum or later.^{26,34,68,75,89,106} Five of these trials were conducted in the USA and one in Germany.⁶⁸ Three trials targeted abstinent pregnant women,^{26,89,106} one of which also included partners,⁸⁹ two trials targeted mixed populations of pregnant smokers and quitters^{34,75} and one trial included postpartum smokers and quitters.⁶⁸ Four out of six interventions included an element of one-to-one counselling plus the provision of self-help materials, whilst two interventions^{26,89} provided only posted self-help educational materials. Control groups received a less intensive intervention or usual care. All long-term effective interventions took place in home settings and were largely delivered remotely (telephone calls, posted or emailed materials) using a variety of intensities and durations. Where counselling was incorporated into the intervention, this was provided by specialist smoking cessation trained

staff or by general trained counsellors. All but one of these trials⁶⁸ reported some biochemical validation of outcome. Two long-term effective trials^{68,75} reported intervention fidelity and/or acceptability concerns: low adherence to motivational interviewing by trained counsellors⁶⁸; or low levels of engaging women in smoking related issues in counselling calls.⁷⁵ Had these issues not been present, it is possible that the intervention effect may have been stronger.

Behaviour change techniques in trials and those characterised as 'promising'

BCTs targeting improvements to smoking behaviour, that we identified from intervention descriptions in our included trials, are summarised in Table 1. BCTs for each individual trial are shown in Supplementary Table 1 (definitions of BCT codes are in Table 1). A total of 45 of the 93 BCTs from the BCTTv1 were identified. The BCTTv1 organises 93 individual BCTs into 16 different groups according to similarity of individual BCT components.¹⁹ All groups were present in our included trials, with the exception of the group related to 'scheduled consequences' (concerning punishments and costs versus rewards). The number of BCTs identified in individual interventions ranged from one to 19, with an average of 7.7 BCTs per intervention. Focusing only on long-term effective interventions,^{26,34,68,75,89,106} the number of BCTs identified ranged from five to 19 with an average of 11 BCTs per intervention. The most prevalent BCTs in all included trials (coded in $\geq 25\%$ interventions) were: 'problem solving' (n=27); 'information about social and environmental consequences' (n=27); 'social support (unspecified)' (n=25); 'information about health consequences' (n=18); 'credible source' (n=17); 'instruction on how to perform a behaviour' (n=11); and 'reduce negative emotions' (n=9). For example, many trials gave information to participants on the health effects of smoking, or effects of exposure to environmental tobacco smoke.

We characterised six BCTs as ‘promising’, or most likely to enhance effectiveness of interventions to maintain smoking abstinence. The BCTs which we identified as ‘promising’ were: ‘problem solving’ in all (i.e., 100%) of long-term effective interventions); ‘information about health consequences’ (100%); ‘information about social and environmental consequences’ (100%); ‘social support’ (100%); ‘reduce negative emotions’ (50%); and ‘instruction on how to perform a behaviour’ (33%). If we were to remove the two trials^{68,75} which reported fidelity and/or acceptability concerns in our analysis, these six BCTs would still meet our definition of ‘promising’, or most likely to be effective.

Discussion

This review is the first study, to our knowledge, to identify BCTs most promising for preventing postpartum smoking relapse. We identified a wide range of BCTs, finding an average of 7.7 BCTs per intervention for all included trials, and an average of 11 BCTs per intervention for long-term effective trials. There is a possibility that the effectiveness of interventions might be improved by incorporating more BCTs, implying more ‘complex’ interventions. We found six ‘promising’ BCTs, which were both frequently occurring and present in interventions that demonstrated long-term effectiveness of sustained smoking abstinence postpartum. These were: ‘problem solving’ which included advice on smoking relapse prevention; ‘information about health consequences’; ‘information about social and environmental consequences’ such as the effects of second-hand smoke; ‘social support’ from a partner or other supporter; ‘reduce negative emotions’ including stress management; and ‘instruction on how to perform a behaviour’ including skills training. Previous studies have identified effective BCTs using the smoking specific taxonomy²² for smoking cessation in pregnancy²⁴; and cessation for smokers in general.^{21,22} This taxonomy has largely been

superseded by the BCTTv1 designed for use in all behaviour change interventions.¹⁹ More recently, Campbell²³ identified BCTs for smoking cessation in pregnancy based on combining the smoking specific taxonomy with the BCTTv1. However, they only included trials from a single Cochrane review.² Our results concur with those of Campbell et al in finding the following BCTs to be potentially effective, based on their presence in two or more effective interventions: ‘information about consequences’; ‘problem solving’; ‘social support’ and ‘reduce negative emotions’. In total their review found 23 potentially effective techniques, but did not find the BCT ‘instruction on how to perform a behaviour’ to be present in effective interventions. However, providing advice on stop smoking medication and changing routine, have been identified as important BCTs for general smokers (smokers not necessarily pregnant or postpartum).^{21,22} Differences in our findings may be since these previous reviews were focused on smoking cessation, whilst we aimed to find BCTs effective for relapse prevention in the postpartum period.

Pregnant and postpartum smokers tend to be of a lower socioeconomic status, less well educated and have lower levels of support.^{2,8} Future interventions and their BCT components need to be tailored to address the specific needs and cultural identity of these high-risk groups. Women who return to smoking postpartum may be less aware of the detrimental effects to the health of the baby caused by their return to smoking.⁸ This suggests the BCTs we identified concerning provision of ‘information about health consequences’ and ‘information about social and environmental consequences’, would be fundamental to the success of an intervention to prevent postpartum smoking relapse. Having greater social or partner support is known to be a key facilitator to prevent relapse^{3,4,8} and our finding of the importance in providing enhanced social support is consistent with this. Only four trials included in this review specifically engaged with partner groups. A review of postpartum relapse strategies concluded that more programmes should include a woman’s partner and her

wider social support network.¹⁰ We found BCTs prompting problem solving and providing guidance on how to maintain smoking abstinence to be ‘promising’, corresponding well with literature finding predictors for relapse include having a lower confidence to remain abstinent postpartum.⁸ Other predictors for relapse are experiencing higher levels of stress, depression and anxiety, and women who smoke are more likely to use substances in response to these circumstances.^{2,8} Our finding of the importance in including the BCT to ‘reduce negative emotions’ might therefore improve the chances of success of an intervention to maintain smoking abstinence.

Many factors influence the effectiveness of BCTs within interventions in addition to the content; including intervention fidelity, acceptability, the mode of delivery, setting, duration, intensity and characteristics of the provider.^{24,131} Considering all of our included trials, we found no clear pattern relating these characteristics to the effectiveness of BCTs although, as noted by others, descriptions of intervention characteristics and fidelity related issues were limited in published papers.¹³¹ Long-term effective interventions that we identified all provided self-help support, and four out of six long-term effective trials combined self-help with counselling. For continued smoking abstinence postpartum, Chamberlain² found some evidence for positive effects of health education, counselling and incentives. For smoking cessation in pregnancy, self-help interventions, including those delivered digitally (by computer or as text messages), have been found to be effective.^{132,133} Self-help provided as written or electronic support, or as part of counselling, may be favourable modes of delivery for BCTs and worthy of further research. Long-term effective interventions included in this study were delivered in home settings and mostly delivered remotely (telephone calls, posted or emailed materials). This suggests effective interventions might be delivered relatively cost-effectively. However, fidelity issues relating to protocol adherence, poor acceptability or participant compliance were evident in two of these trials,^{68,75} which may have limited

intervention effectiveness. Future interventions need to consider issues of implementation and acceptability in this population, who may be challenging to engage.

Potential limitations to this review included the heterogeneity of participants, types of interventions and outcomes. Therefore, as in similar BCT content reviews,^{23,24} we did not attempt meta-analysis. As noted elsewhere in the literature,¹³¹ the level of detail needed for BCT coding was often not present in published intervention descriptions. In practice, interventions may have used more BCTs than those reported. We did not contact study authors, but did address this by taking an inclusive approach in our coding, including BCTs coded as probably present (+) in addition to those coded as definitely (++) present (www.bct-taxonomy.com). We included a second independent coder for 10% of data extraction and involved a third reviewer where necessary to resolve discrepancies and to ensure any relevant BCTs had been correctly identified. Unfortunately we were unable to fully duplicate screening and data extraction due to limited resources. We did not assess risk of bias, as our main aim was to identify any potentially relevant BCTs, but we did describe any reported fidelity and/or acceptability issues, to consider how BCTs might be best delivered. As in other reviews²³ we did not compare BCTs within effective and non-effective interventions. No trials reported a negative intervention effect and using this approach may have limited the number of potentially relevant BCTs. Measures of study effectiveness are also acknowledged to be ambiguous, since statistical significance is dependent on a wide range of factors, such as sample size.¹³⁴ We cannot say with certainty that the presence of individual BCTs, demonstrates any causal relationship with trial outcome, but their repeated presence across long-term effective trials suggests they may be more promising approaches for future intervention content.

Not all trials validated smoking abstinence biochemically, although a Cochrane review reported similar effect sizes in pregnancy trials which were validated and trials which were not.² All trials were conducted in high-income countries with the majority conducted in the USA; which may mean findings would not be generalizable to other countries or contexts. Smoking in pregnancy is reducing in higher income countries but increasing in low and middle-income countries^{2,16} and no such countries met the inclusion criteria for this review. However, this may be since we excluded articles not published in English. The majority of included trials were targeted to general populations of pregnant or postpartum women, rather than populations of ethnic minorities or lower socioeconomic groups, where smoking prevalence is higher. Further research with these specific groups would be beneficial and might identify further relevant BCTs, such as BCTs concerning identity specific to different populations. Difficulty in adapting to a new mothering identity has been identified as a key factor in prompting return to smoking postpartum.^{3,4} We did not identify any trials that focused on nicotine replacement therapy (NRT) or were recent enough to include more novel approaches such as e-cigarettes. Evidence suggests that e-cigarettes are significantly less harmful than smoking tobacco and many smokers use e-cigarettes as an aid to quit.^{1,135} There is some evidence that for pharmacological interventions for smoking cessation in pregnancy, behavioural support in combination with NRT might be helpful.¹⁶

Strengths to this review included the systematic approach to identifying relevant interventions and the application of the most recent behaviour change technique taxonomy.¹⁹ We characterised BCTs as ‘promising’ if present in long-term effective interventions and where frequency in all included trials (regardless of effectiveness) was high. We were therefore able to consider ‘promising’ BCTs in light of their overall frequency in interventions relevant to this population, giving an indication of the ease by which these BCTs might be incorporated into interventions, their suitability to the setting, feasibility and

acceptability. This is one of the first systematic reviews of BCTs specifically to prevent return to smoking postpartum and these results further our understanding of what behavioural support might be effective. Future research could compare interventions with different combinations of BCTs, or could compare BCTs in intervention and control groups. The identification and testing of BCTs in future postpartum relapse prevention interventions in a more diverse range of countries, with more diverse populations, and incorporating more novel components such as e-cigarettes would be beneficial to research and future practice. Identifying six specific BCTs as promising components for postpartum smoking relapse prevention highlights the need to customise support for this particular population. Further research by our team will explore these BCT components through qualitative research with postpartum ex-smokers in order to develop a prototype intervention for maintenance of smoking abstinence postpartum (MRC PHIND grant ref: MR/P016944/1).

Conclusions

We identified six promising BCTs to prevent postpartum smoking relapse based on a structured systematic review of published available evidence. Future interventions should consider the inclusion of BCTs addressing problem solving and how to maintain abstinent behaviour, information on health and other consequences of smoking, reducing negative emotions and improving the likelihood of smoking abstinence through the provision of social support.

Funding

This work was supported by the Medical Research Council (PHIND grant ref: MR/P016944/1).

Declaration of Interests

None declared.

Acknowledgements

With thanks to Isabel Greaves (IG) for her assistance in screening articles for inclusion in this review.

References

1. DoH. Towards a Smokefree Generation. A Tobacco Control Plan for England. London: Department of Health; 2017.
2. Chamberlain C, O'Mara-Eves A, Porter J, et al. Psychosocial interventions for supporting women to stop smoking in pregnancy. Cochrane Database Syst Rev. 2017(2):Cd001055. DOI: 001010.001002/14651858.CD14001055.pub14651855.
3. Flemming K, McCaughan D, Angus K, Graham H. Qualitative systematic review: barriers and facilitators to smoking cessation experienced by women in pregnancy and following childbirth. J Adv Nurs. 2015;71(6):1210-1226.
4. Notley C, Blyth A, Craig J, Edwards A, Holland R. Postpartum smoking relapse-a thematic synthesis of qualitative studies. Addiction. 2015;110(11):1712-1723.
5. NICE. How to Stop Smoking in Pregnancy and Following Childbirth: Public Health Guideline [PH26]. London: National Institute for Health and Clinical Excellence;2010.

6. Jones M, Lewis S, Parrott S, Wormald S, Coleman T. Re-starting smoking in the postpartum period after receiving a smoking cessation intervention: a systematic review. *Addiction*. 2016;111(6):981-990.
7. Lumley J, Chamberlain C, Dowswell T, Oliver S, Oakley L, Watson L. Interventions for promoting smoking cessation during pregnancy. *Cochrane Database Syst Rev*. 2009(3):Cd001055. DOI: 001010.001002/14651858.CD14001055.pub14651853.
8. Orton S, Coleman T, Coleman-Haynes T, Ussher M. Predictors of postpartum return to smoking: a systematic review. *Nicotine Tob Res*. 2018;20(6):665-673.
9. Riaz M, Lewis S, Naughton F, Ussher M. Predictors of smoking cessation during pregnancy: a systematic review and meta-analysis. *Addiction*. 2018;113(4):610-622.
10. Fang WL, Goldstein AO, Butzen AY, et al. Smoking cessation in pregnancy: a review of postpartum relapse prevention strategies. *J Am Board Fam Pract*. 2004;17(4):264-275.
11. WHO. WHO Recommendations for the Prevention and Management of Tobacco Use and Second-Hand Smoke Exposure in Pregnancy. Geneva, Switzerland: World Health Organization; 2013.
12. Godfrey C, Pickett K, Parrott S, Mdege N, Eapen D. Estimating the Costs to the NHS of Smoking in Pregnancy for Pregnant Women and Infants. York: Public Health Research Consortium;2010.
13. Pirie K, Peto R, Reeves GK, Green J, Beral V. The 21st century hazards of smoking and benefits of stopping: a prospective study of one million women in the UK. *Lancet*. 2013;381(9861):133-141.
14. RCP. Passive Smoking and Children. London: Royal College of Physicians;2010.

15. Leonardi-Bee J, Jere ML, Britton J. Exposure to parental and sibling smoking and the risk of smoking uptake in childhood and adolescence: a systematic review and meta-analysis. *Thorax*. 2011;66(10):847-855.
16. Coleman T, Chamberlain C, Davey M-A, Cooper SE, Leonardi-Bee J. Pharmacological interventions for promoting smoking cessation during pregnancy. *Cochrane Database Syst Rev*. 2015(12):Cd010078. DOI: 010010.011002/14651858.CD14010078.pub14651852.
17. Hajek P, Stead LF, West R, Jarvis M, Hartmann-Boyce J, Lancaster T. Relapse prevention interventions for smoking cessation. *Cochrane Database Syst Rev*. 2013(8):Cd003999. DOI: 003910.001002/14651858.CD14003999.pub14651854.
18. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ*. 2008;337:a1655.
19. Michie S, Richardson M, Johnston M, et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med*. 2013;46(1):81-95.
20. Michie S, Wood CE, Johnston M, Abraham C, Francis JJ, Hardeman W. Behaviour change techniques: the development and evaluation of a taxonomic method for reporting and describing behaviour change interventions (a suite of five studies involving consensus methods, randomised controlled trials and analysis of qualitative data). *Health Technol Assess*. 2015;19(99):1-188.
21. Lorencatto F, West R, Seymour N, Michie S. Developing a method for specifying the components of behavior change interventions in practice: the example of smoking cessation. *J Consult Clin Psychol*. 2013;81(3):528-544.

22. West R, Walia A, Hyder N, Shahab L, Michie S. Behavior change techniques used by the English Stop Smoking Services and their associations with short-term quit outcomes. *Nicotine Tob Res.* 2010;12(7):742-747.
23. Campbell KA, Fergie L, Coleman-Haynes T, et al. Improving behavioral support for smoking cessation in pregnancy: what are the barriers to stopping and which behavior change techniques can influence these? Application of Theoretical Domains Framework. *Int J Environ Res Public Health.* 2018;15(2):DOI: 10.3390/ijerph15020359.
24. Lorencatto F, West R, Michie S. Specifying evidence-based behavior change techniques to aid smoking cessation in pregnancy. *Nicotine Tob Res.* 2012;14(9):1019-1026.
25. Allen SS, Allen AM, Lunos S, Tosun N. Progesterone and postpartum smoking relapse: A pilot double-blind placebo-controlled randomized trial. *Nicotine Tob Res.* 2016;18(11):2145-2153.
26. Brandon TH, Simmons VN, Meade CD, et al. Self-help booklets for preventing postpartum smoking relapse: a randomized trial. *Am J Public Health.* 2012;102(11):2109-2115.
27. Correa JB, Simmons VN, Sutton SK, Meltzer LR, Brandon TH. A content analysis of attributions for resuming smoking or maintaining abstinence in the post-partum period. *Matern Child Health J.* 2015;19(3):664-674.
28. Lopez ES, M. Meade, C. Quinn, G. Pedraza, J. Brandon, T. The elusive pregnant ex-smoker: lessons from recruitment for a clinical trial. Society for Research on Nicotine and Tobacco 11th Annual Meeting, Prague. 2005.

29. Lopez EN, Simmons VN, Quinn GP, Meade CD, Chirikos TN, Brandon TH. Clinical trials and tribulations: lessons learned from recruiting pregnant ex-smokers for relapse prevention. *Nicotine Tob Res.* 2008;10(1):87-96.
30. Quinn G, Ellison BB, Meade C, et al. Adapting smoking relapse-prevention materials for pregnant and postpartum women: formative research. *Matern Child Health J.* 2006;10(3):235-245.
31. Simmons VN, Lopez Khoury E, Segall Koltz EJ, et al. Preventing smoking relapse among pregnant and postpartum women: a randomized clinical trial. Joint conference of SRNT and SRNT-E, Dublin. 2009:POS2-16.
32. Simmons VN, Cruz LM, Brandon TH, Quinn GP. Translation and adaptation of smoking relapse-prevention materials for pregnant and postpartum Hispanic women. *J Health Commun.* 2011;16(1):90-107.
33. Simmons VN, Sutton SK, Quinn GP, Meade CD, Brandon TH. Prepartum and postpartum predictors of smoking. *Nicotine Tob Res.* 2014;16(4):461-468.
34. Cummins SE, Tedeschi GJ, Anderson CM, Zhu SH. Telephone intervention for pregnant smokers: a randomized controlled trial. *Am J Prev Med.* 2016;51(3):318-326.
35. Cummins SE, Tedeschi GJ, Anderson CM, Quinlan-Downs R, Harris P, Zhu S-H. Telephone counselling for pregnant smokers: essential elements. *J Smok Cessat.* 2007;2(2):36-46.
36. El-Mohandes AA, El-Khorazaty MN, Kiely M, Gantz MG. Smoking cessation and relapse among pregnant African-American smokers in Washington, DC. *Matern Child Health J.* 2011;15 Suppl 1:S96-105.
37. Backonja U, Robledo CA, Wallace ME, Flores KF, Kiely M. Reproductive health knowledge among African American women enrolled in a clinic-based randomized

- controlled trial to reduce psychosocial and behavioral risk: Project DC-HOPE. *Women's Health Issues*. 2016;26(4):442-451.
38. Blake S, Joseph J, Schwartz D, El-Khorazaty N, Gantz M, El-Mohandes A. Preventing prenatal and postpartum environmental tobacco smoke (ETS) exposure. *Pediatric Academic Societies Annual Meeting*; 2005; Washington, DC.
 39. Blake SE-M, A. Schwartz, D. El-Khorazaty, N. Gantz, M. Joseph, J. Promoting smoking cessation during pregnancy and preventing postpartum relapse. *Pediatric Academic Societies Annual Meeting*; 2005; Washington, DC.
 40. Blake SM, Murray KD, El-Khorazaty MN, et al. Environmental tobacco smoke avoidance among pregnant African-American nonsmokers. *Am J Prev Med*. 2009;36(3):225-234.
 41. El-Khorazaty MN, Johnson AA, Kiely M, et al. Recruitment and retention of low-income minority women in a behavioral intervention to reduce smoking, depression, and intimate partner violence during pregnancy. *BMC Public Health*. 2007;7:233.
 42. El-Mohandes AAE. An integrated psycho-behavioral intervention during pregnancy has significant effects in reducing risks during the post-partum period in African-American women. *Pediatric Academic Societies Annual Meeting*; 2005; Washington, DC.
 43. El-Mohandes AAE. A psycho-behavioral intervention on African American pregnant women with a history of intimate partner violence (IPV) improves birth weight distribution of their newborns. *Pediatric Academic Societies Annual Meeting*; 2005; San Francisco, CA.
 44. El-Mohandes AE. An integrated behavioral intervention reduces rates of moderate and extreme prematurity in African American (AA) mothers with a history of

- smoking during pregnancy. Pediatric Academic Societies Annual Meeting; 2006; San Francisco, CA.
45. El-Mohandes AAE, Kiely M, Gantz MG, El-Khorazaty N. A multiple risk factor behavioral intervention reduces environmental tobacco smoke exposure. Pediatric Academic Societies Annual Meeting; 2007; Toronto, Canada.
 46. El-Mohandes AAE, Kiely M, Joseph JG, et al. An integrated intervention in pregnant African Americans reduces postpartum risk: a randomized trial. *Obstet Gynecol.* 2008;112(3):611-620.
 47. El-Mohandes AA, Kiely M, Blake SM, Gantz MG, El-Khorazaty MN. An intervention to reduce environmental tobacco smoke exposure improves pregnancy outcomes. *Pediatrics.* 2010;125(4):721-728.
 48. El-Mohandes AAE, Kiely M, Gantz MG, El-Khorazaty MN. Very preterm birth is reduced in women receiving an integrated behavioral intervention: a randomized controlled trial. *Matern Child Health J.* 2011;15(1):19-28.
 49. Joseph J. Randomized trial to reduce 4 behaviors linked to adverse pregnancy outcomes among 1048 inner-city African American women. Pediatric Academic Societies Annual Meeting; 2005; Washington, DC.
 50. Joseph JG, El-Mohandes AA, Kiely M, et al. Reducing psychosocial and behavioral risk factors: results of a multisite intervention randomized trial of high-risk pregnant African American women. *Am J Pub Health.* 2009;99(6):1053-1061.
 51. Katz KS, Blake SM, Milligan RA, et al. The design, implementation and acceptability of an integrated intervention to address multiple behavioral and psychosocial risk factors among pregnant African American women. *BMC Pregnancy Childbirth.* 2008;8:22.

52. Kiely M, El-Khorazaty MN, El-Mohandes AAE. Depression and smoking during pregnancy impact the efficacy of an integral behavioral intervention to resolve risks. Pediatric Academic Societies Annual Meeting; 2007; Toronto, Canada.
53. Subramanian S, Katz KS, Rodan M, et al. An integrated randomized intervention to reduce behavioral and psychosocial risks: pregnancy and neonatal outcomes. *Matern Child Health J.* 2012;16(3):545-554.
54. Tan S, Courtney LP, El-Mohandes AA, et al. Relationships between self-reported smoking, household environmental tobacco smoke exposure and depressive symptoms in a pregnant minority population. *Matern Child Health J.* 2011;15 Suppl 1:S65-74.
55. Edwards NC, Sims-Jones N. A randomized controlled trial of alternative approaches to community follow-up for postpartum women. *Can J Public Health.* 1997;88(2):123-128.
56. Edwards N, Sims-Jones N. Smoking and smoking relapse during pregnancy and postpartum: results of a qualitative study. *Birth.* 1998;25(2):94-100.
57. Ershoff DH, Quinn VP, Mullen PD. Relapse prevention among women who stop smoking early in pregnancy: a randomized clinical trial of a self-help intervention. *Am J Prev Med.* 1995;11(3):178-184.
58. Ershoff DH, Mullen PD, Quinn VP. A randomized trial of a serialized self-help smoking cessation program for pregnant women in an HMO. *Am J Pub Health.* 1989;79(2):182-187.
59. Ershoff DH, Quinn VP, Mullen PD, Lairson DR. Pregnancy and medical cost outcomes of a self-help prenatal smoking cessation program in a HMO. *Public Health Rep.* 1990;105(4):340-347.

60. Mullen PD, Quinn VP, Ershoff DH. Maintenance of nonsmoking postpartum by women who stopped smoking during pregnancy. *Am J Public Health.* 1990;80(8):992-994.
61. Mullen PD, Carbonari JP, Tabak ER, Glenday MC. Improving disclosure of smoking by pregnant women. *Am J Obstet Gynecol.* 1991;165(2):409-413.
62. Mullen PD, Richardson MA, Quinn VP, Ershoff DH. Postpartum return to smoking: who is at risk and when. *Am J Health Promot.* 1997;11(5):323-330.
63. Quinn VP, Mullen PD, Ershoff DH. Women who stop smoking spontaneously prior to prenatal care and predictors of relapse before delivery. *Addict Behav.* 1991;16(1-2):29-40.
64. Forray A, Waters A. Ecological momentary assessments and attentional bias modification for postpartum smoking. *Neuropsychopharmacology.* 2015;40:S590-S591.
65. Foster DW, Waters AJ, Forray A. Perinatal attentional retraining intervention for smoking - A pilot study. *Drug Alcohol Depend.* 2015;156:e73.
66. Forray A, Gilstad-Hayden K, Sofuoglu M, Yonkers K. Progesterone for postpartum smoking relapse prevention. *Neuropsychopharmacology.* 2016;41:S606.
67. Hajek P, West R, Lee A, et al. Randomized controlled trial of a midwife-delivered brief smoking cessation intervention in pregnancy. *Addiction.* 2001;96(3):485-494.
68. Hannover W, Thyrian JR, Roske K, et al. Smoking cessation and relapse prevention for postpartum women: results from a randomized controlled trial at 6, 12, 18 and 24 months. *Addict Behav.* 2009;34(1):1-8.
69. Roske K, Hannover W, Grempler J, et al. Post-partum intention to resume smoking. *Health Educ Res.* 2006;21(3):386-392.

70. Roske K, Schumann A, Hannover W, et al. Postpartum smoking cessation and relapse prevention intervention: a structural equation modeling application to behavioral and non-behavioral outcomes of a randomized controlled trial. *J Health Psychol.* 2008;13(4):556-568.
71. Thyrian JR, Hannover W, Grempler J, Roske K, John U, Hapke U. An intervention to support postpartum women to quit smoking or remain smoke-free. *J Midwifery Womens Health.* 2006;51(1):45-50.
72. Thyrian JR, Hannover W, Roske K, Rumpf HJ, John U, Hapke U. Postpartum return to smoking: identifying different groups to tailor interventions. *Addict Behav.* 2006;31(10):1785-1796.
73. Thyrian JR, Freyer-Adam J, Hannover W, et al. Adherence to the principles of motivational interviewing, clients' characteristics and behavior outcome in a smoking cessation and relapse prevention trial in women postpartum. *Addict Behav.* 2007;32(10):2297-2303.
74. Thyrian JR, Freyer-Adam J, Hannover W, et al. Population-based smoking cessation in women post partum: adherence to motivational interviewing in relation to client characteristics and behavioural outcomes. *Midwifery.* 2010;26(2):202-210.
75. McBride CM, Curry SJ, Lando HA, Pirie PL, Grothaus LC, Nelson JC. Prevention of relapse in women who quit smoking during pregnancy. *Am J Public Health.* 1999;89(5):706-711.
76. Curry SJ, McBride C, Grothaus L, Lando H, Pirie P. Motivation for smoking cessation among pregnant women. *Psychol Addict Behav.* 2001;15(2):126-132.
77. Lando HA, Valanis BG, Lichtenstein E, et al. Promoting smoking abstinence in pregnant and postpartum patients: a comparison of 2 approaches. *Am J Manag Care.* 2001;7(7):685-693.

78. McBride CM, Curry SJ, Grothaus LC, Nelson JC, Lando H, Pirie PL. Partner smoking status and pregnant smoker's perceptions of support for and likelihood of smoking cessation. *Health Psychol.* 1998;17(1):63-69.
79. Jimenez-Muro A, Nerin I, Samper P, et al. A proactive smoking cessation intervention in postpartum women. *Midwifery.* 2013;29(3):240-245.
80. Johnson JL, Ratner PA, Bottorff JL, Hall W, Dahinten S. Preventing smoking relapse in postpartum women. *Nurs Res.* 2000;49(1):44-52.
81. Bottorff JL, Johnson JL, Irwin LG, Ratner PA. Narratives of smoking relapse: the stories of postpartum women. *Res Nurs Health.* 2000;23(2):126-134.
82. Ratner PA, Johnson JL, Bottorff JL. Smoking relapse and early weaning among postpartum women: is there an association? *Birth.* 1999;26(2):76-82.
83. Kientz E, Kupperschmidt B. KICCS: a successful strategy to promote smoking cessation in women during and post pregnancy. *Okla Nurse.* 2005;50(4):27-30.
84. Lillington L, Royce J, Novak D, Ruvalcaba M, Chlebowski R. Evaluation of a smoking cessation program for pregnant minority women. *Cancer Pract.* 1995;3(3):157-163.
85. McBride CM, Baucom DH, Peterson BL, et al. Prenatal and postpartum smoking abstinence a partner-assisted approach. *Am J Prev Med.* 2004;27(3):232-238.
86. Morasco BJ, Dornelas EA, Fischer EH, Oncken C, Lando HA. Spontaneous smoking cessation during pregnancy among ethnic minority women: a preliminary investigation. *Addict Behav.* 2006;31(2):203-210.
87. Dornelas EA, Magnavita J, Beazoglou T, et al. Efficacy and cost-effectiveness of a clinic-based counseling intervention tested in an ethnically diverse sample of pregnant smokers. *Patient Educ Couns.* 2006;64(1-3):342-349.

88. Winickoff JP, Healey EA, Regan S, et al. Using the postpartum hospital stay to address mothers' and fathers' smoking: The NEWS Study. *Pediatrics*. 2010;125(3):518.
89. Mullen PD, DiClemente CC, Bartholomew LK. Theory and context in project PANDA: a program to help postpartum women stay off cigarettes. In: Bartholomew LK, Parcel GS, Kok G, Gottlieb NH, eds. *Intervention Mapping: A Process for Designing Theory- and Evidence-based Health Promotion Programs*. Palo Alto, CA: Mayfield: McGraw-Hill Publishing Co.; 2001.
90. DiClemente C, Dolan-Mullen P, Windsor R. The process of pregnancy smoking cessation: implications for interventions. *Tob Control*. 2000;9(Suppl 3):iii16-iii21.
91. Mullen PD, DiClemente C, Carbonari J, et al. Project PANDA maintenance of pre-natal smoking abstinence postpartum at 6 weeks and 3, 6, and 12 months. *Ann Behav Med*. 1997;19:130.
92. Sockrider MM, Hudmon KS, Addy R, Dolan Mullen P. An exploratory study of control of smoking in the home to reduce infant exposure to environmental tobacco smoke. *Nicotine Tob Res*. 2003;5(6):901-910.
93. Stotts AL, DiClemente CC, Carbonari JP, Mullen PD. Postpartum return to smoking: staging a "suspended" behavior. *Health Psychol*. 2000;19(4):324-332.
94. Petersen L, Handel J, Kotch J, Podedworny T, Rosen A. Smoking reduction during pregnancy by a program of self-help and clinical support. *Obstet Gynecol*. 1992;79:924-930.
95. Polanska K, Hanke W, Sobala W, Lowe JB. Efficacy and effectiveness of the smoking cessation program for pregnant women. *Int J Occup Med Environ Health*. 2004;17(3):369-377.

96. Polanska K, Hanke W, Sobala W. Smoking relapse one year after delivery among women who quit smoking during pregnancy. *Int J Occup Med Environ Health*. 2005;18(2):159-165.
97. Polanska K, Hanke W, Sobala W. Characteristic of the smoking habit among pregnant women on the base of the test "Why am I smoker?". *Przegl Lek*. 2005;62(10):1095-1098.
98. Strecher VJ, Bishop KR, Bernhardt J, Thorp JM, Cheuvront B, Potts P. Quit for keeps: tailored smoking cessation guides for pregnancy and beyond. *Tob Control*. 2000;9 Suppl 3:iii78-79.
99. Pollak KI, Fish LJ, Lyna P, et al. Efficacy of a nurse-delivered intervention to prevent and delay postpartum return to smoking: The quit for two trial. *Nicotine Tob Res*. 2016;18(10):1960-1966.
100. Pollak KI, Fish LJ, Lyna P, Peterson BL, Swamy GK, Levine MD. Predictors of pregnant quitters' intention to return to smoking postpartum. *Nicotine Tob Res*. 2015;17(6):742-745.
101. Pbert L, Ockene JK, Zapka J, et al. A community health center smoking-cessation intervention for pregnant and postpartum women. *Am J Prev Med*. 2004;26(5):377-385.
102. Bonollo DP, Zapka JG, Stoddard AM, Ma Y, Pbert L, Ockene JK. Treating nicotine dependence during pregnancy and postpartum: understanding clinician knowledge and performance. *Patient Educ Couns*. 2002;48(3):265-274.
103. Ma Y, Goins KV, Pbert L, Ockene JK. Predictors of smoking cessation in pregnancy and maintenance postpartum in low-income women. *Matern Child Health J*. 2005;9(4):393-402.

104. Zapka JG, Pbert L, Stoddard AM, Ockene JK, Goins KV, Bonollo D. Smoking cessation counseling with pregnant and postpartum women: a survey of community health center providers. *Am J Public Health*. 2000;90(1):78-84.
105. Zapka J, Goins KV, Pbert L, Ockene JK. Translating efficacy research to effectiveness studies in practice: lessons from research to promote smoking cessation in community health centers. *Health Promot Pract*. 2004;5(3):245-255.
106. Reitzel LR, Vidrine JJ, Businelle MS, et al. Preventing postpartum smoking relapse among diverse low-income women: a randomized clinical trial. *Nicotine Tob Res*. 2010;12(4):326-335.
107. Businelle MS, Kendzor DE, Reitzel LR, et al. Pathways linking socioeconomic status and postpartum smoking relapse. *Ann Behav Med*. 2013;45(2):180-191.
108. Correa-Fernandez V, Ji L, Castro Y, et al. Mediators of the association of major depressive syndrome and anxiety syndrome with postpartum smoking relapse. *J Consult Clin Psychol*. 2012;80(4):636-648.
109. Heppner WL, Ji L, Reitzel LR, et al. The role of prepartum motivation in the maintenance of postpartum smoking abstinence. *Health Psychol*. 2011;30(6):736-745.
110. Kendzor DE, Businelle MS, Costello TJ, et al. Breast feeding is associated with postpartum smoking abstinence among women who quit smoking due to pregnancy. *Nicotine Tob Res*. 2010;12(10):983-988.
111. Ruger JP, Weinstein MC, Hammond SK, Kearney MH, Emmons KM. Cost-effectiveness of motivational interviewing for smoking cessation and relapse prevention among low-income pregnant women: a randomized controlled trial. *Value Health*. 2008;11(2):191-198.

112. Ruger JP, Emmons KM, Kearney MH, Weinstein MC. Measuring the costs of outreach motivational interviewing for smoking cessation and relapse prevention among low-income pregnant women. *BMC Pregnancy Childbirth*. 2009;9:46-46.
113. Kendrick JS, Zahniser SC, Miller N, et al. Integrating smoking cessation into routine public prenatal care: the Smoking Cessation in Pregnancy project. *Am J Public Health*. 1995;85(2):217-222.
114. England LJ, Kendrick JS, Wilson HG, Merritt RK, Gargiullo PM, Zahniser SC. Effects of smoking reduction during pregnancy on the birth weight of term infants. *Am J Epidemiol*. 2001;154(8):694-701.
115. Spierto FW, Hannon W, Kendrick J, Bernert J, Pirkle J, Pm G. Urinary cotinine levels in women enrolled in a smoking cessation study during and after pregnancy. *J Smoking-Related Dis*. 1994;5(2):65-76.
116. Secker-Walker RH, Solomon LJ, Flynn BS, et al. Smoking relapse prevention counseling during prenatal and early postnatal care. *Am J Prev Med*. 1995;11(2):86-93.
117. Secker-Walker RH, Solomon LJ, Flynn BS, et al. Individualized smoking cessation counseling during prenatal and early postnatal care. *Am J Obstet Gynecol*. 1994;171(5):1347-1355.
118. Secker-Walker RH, Solomon LJ, Flynn BS, Skelly JM, Mead PB. Smoking relapse prevention during pregnancy. A trial of coordinated advice from physicians and individual counseling. *Am J Prev Med*. 1998;15(1):25-31.
119. Secker-Walker RH, Solomon LJ, Flynn BS, et al. Training obstetric and family practice residents to give smoking cessation advice during prenatal care. *Am J Obstet Gynecol*. 1992;166(5):1356-1363.

120. Secker-Walker RH, Vacek PM, Flynn BS, Mead PB. Smoking in pregnancy, exhaled carbon monoxide, and birth weight. *Obstet Gynecol.* 1997;89(5 Pt 1):648-653.
121. Secker-Walker RH, Solomon LJ, Flynn BS, Skelly JM, Mead PB. Reducing smoking during pregnancy and postpartum: physician's advice supported by individual counseling. *Prev Med.* 1998;27(3):422-430.
122. Secker-Walker RH, Vacek PM, Flynn BS, Mead PB. Estimated gains in birth weight associated with reductions in smoking during pregnancy. *J Reprod Med.* 1998;43(11):967-974.
123. Solomon LJ, Secker-Walker RH, Skelly JM, Flynn BS. Stages of change in smoking during pregnancy in low-income women. *J Behav Med.* 1996;19(4):350-366.
124. Levine MD, Cheng Y, Marcus MD, Kalarchian MA, Emery RL. Preventing postpartum smoking relapse: A randomized clinical trial. *Jama Intern Med.* 2016;176(4):443-452.
125. Emery RL, Gregory MP, Grace JL, Levine MD. Prevalence and correlates of a lifetime cannabis use disorder among pregnant former tobacco smokers. *Addict Behav.* 2016;54:52-58.
126. Levine MD, Marcus MD, Kalarchian MA, Cheng Y. Strategies to Avoid Returning to Smoking (STARTS): A randomized controlled trial of postpartum smoking relapse prevention interventions. *Contemp Clin Trials.* 2013;36(2):10.1016/j.cct.2013.1010.1002.
127. Levine MD, Cheng Y, Marcus MD, Emery RL. Psychiatric disorders and gestational weight gain among women who quit smoking during pregnancy. *J Psychosom Res.* 2015;78(5):504-508.
128. Suplee PD. The importance of providing smoking relapse counseling during the postpartum hospitalization. *J Obstet Gynecol Neonatal Nurs.* 2005;34(6):703-712.

129. Thornton L. The Rotunda Stop Smoking Programme: Smoking and Pregnancy: Feasability and Effectiveness of a Smoking Intervention Programme Among Pregnant Women. Ireland: Department of Public Health, Eastern Health Board;1997.
130. Thornton L, Gogan C, McKenna P. The rotunda stop smoking programme. *Irish J Med Sci.* 1998;167(Suppl 9):28.
131. Michie S, Fixsen D, Grimshaw JM, Eccles MP. Specifying and reporting complex behaviour change interventions: the need for a scientific method. *Implement Sci.* 2009;4:40.
132. Griffiths SE, Parsons J, Naughton F, Fulton EA, Tombor I, Brown KE. Are digital interventions for smoking cessation in pregnancy effective? A systematic review and meta-analysis. *Health Psychol Rev.* 2018:1-24.
133. Naughton F, Prevost AT, Sutton S. Self-help smoking cessation interventions in pregnancy: a systematic review and meta-analysis. *Addiction.* 2008;103(4):566-579.
134. Peters GJ, de Bruin M, Crutzen R. Everything should be as simple as possible, but no simpler: towards a protocol for accumulating evidence regarding the active content of health behaviour change interventions. *Health Psychol Rev.* 2015;9(1):1-14.
135. McNeill A, Brose LS, Calder R, Bauld L, Robson D. Evidence Review of E-cigarettes and Heated Tobacco Products. London: Public Health England; 2018.

Table 1: Frequency of BCTs identified in smoking relapse interventions

BCT code	BCT label	BCT in all studies; n (%); Max n=32	BCT in 'long-term effective' [#] studies; n (%); Max n=6
1.1	Goal setting (behavior)	5 (16)	1 (17)
1.2	Problem solving	27 (84)	6 (100) *
1.4	Action planning	1 (3)	0 (0)
1.5	Review behavior goal(s)	4 (13)	0 (0)
1.6	Discrepancy between current behavior and goal	1 (3)	1 (17)
1.7	Review outcome goal(s)	2 (6)	0 (0)
1.8	Behavioral contract	2 (6)	0 (0)
1.9	Commitment	3 (9)	1 (17)
2.2	Feedback on behavior	1 (3)	0 (0)
2.3	Self-monitoring of behavior	3 (9)	0 (0)
2.6	Biofeedback	4 (13)	0 (0)
3.1	Social support (unspecified)	25 (78)	6 (100) *
3.2	Social support (practical)	3 (9)	1 (17)
3.3	Social support (emotional)	3 (9)	1 (17)
4.1	Instruction on how to perform a behavior	11 (34)	2 (33) *
4.2	Information about antecedents	2 (6)	0 (0)
4.3	Re-attribution	1 (3)	1 (17)
5.1	Information about health consequences	18 (56)	6 (100) *
5.3	Information about social and environmental consequences	27 (84)	6 (100) *
5.6	Information about emotional consequences	1 (3)	1 (17)
6.1	Demonstration of the behavior	2 (6)	1 (17)
6.2	Social comparison	4 (13)	3 (50)
7.1	Prompts/cues	4 (13)	1 (17)
8.1	Behavioral practice/ rehearsal	2 (6)	0 (0)
8.2	Behavior substitution	5 (16)	2 (33)
9.1	Credible source	17 (53)	1 (17)
9.2	Pros and cons	7 (22)	3 (50)
9.3	Comparative imagining of future outcomes	1 (3)	0 (0)
10.1	Material incentive (behavior)	1 (3)	0 (0)
10.2	Material reward (behavior)	1 (3)	0 (0)
10.3	Non-specific reward	1 (3)	0 (0)
10.4	Social reward	6 (19)	2 (33)
10.7	Self-incentive	1 (3)	0 (0)
11.1	Pharmacological support	3 (9)	1 (17)
11.2	Reduce negative emotions	9 (28)	3 (50) *
12.1	Restructuring the physical environment	2 (6)	1 (17)
12.2	Restructuring the social environment	2 (6)	1 (17)
12.3	Avoidance/reducing exposure to cues for the behavior	4 (13)	2 (33)
12.5	Adding objects to the environment	4 (13)	2 (33)
13.1	Identification of self as role model	3 (9)	2 (33)
13.2	Framing/reframing	7 (22)	2 (33)
13.5	Identity associated with changed behavior	2 (6)	2 (33)
15.1	Verbal persuasion about capability	7 (22)	1 (17)
15.3	Focus on past success	1 (3)	1 (17)
16.2	Imaginary reward	3 (9)	2 (33)

[#] p<0.05 in intervention vs control at ≥ 6 months postpartum; *BCTs identified as 'promising'

ACCEPTED MANUSCRIPT

Highlights:

- First review of behaviour change techniques to prevent postpartum smoking relapse
- Six promising behaviour change techniques (BCTs) were frequently coded
- BCTs were problem solving, social support, information about consequences
- How to perform a behaviour and reduce negative emotions were also promising BCTs




Figure 1